

Proxy Simulation Schemes

for Generic Robust Monte-Carlo Sensitivities and High Accuracy Drift Approximation

with Applications to the LIBOR Market Model

Christian Fries
16.12.2005
(Version 1.1)

www.christian-fries.de/finmath/talks/2005qmf

Agenda

- Monte Carlo Methods
 - Temporal Discretisation Error
 - Sensitivity Calculation
 - Finite Differences
 - Pathwise Differentiation
 - Pathwise Differentiation (alternative view)
 - Likelihood Ratio Method
 - Malliavin Calculus
- Proxy Simulations Scheme Method
 - Pricing & Sensitivity Calculation
 - Densities & Weak Schemes
 - Implementation
- Example: LIBOR Market Model
- Properties
 - Degenerate Diffusion Matrix / Measure Equivalence
- Numerical Results
 - Drift Approximations
 - Sensitivity Calculations
- Appendix
 - A Quadratic WKB Expansion for the LMM Transition Probability
- References

Monte Carlo Methods

Challenges

Discretization Error:

Drift Approximations

Monte Carlo Methods: Discretization Error

Consider for SDE

$$dX(t) = \mu(t, X(t))X(t)dt + \sigma(t, X(t))X(t)dW(t),$$

e.g. the Log-Euler Scheme

$$X(t + \Delta t) = X(t) \cdot \exp \left(\mu(t, X(t))\Delta t - \frac{1}{2}\sigma^2(t, X(t))\Delta t + \sigma(t, X(t))\Delta W(t) \right).$$

If σ is constant on $[t, \Delta t]$ (Black Model, LIBOR Market Model) but μ is stochastic and/or non-linear (LIBOR Market Model), then the discretization error is given by a drift approximation error, e.g. here

$$\int_t^{t+\Delta t} \mu(\tau, X(\tau))d\tau \approx \mu(t, X(t))\Delta t.$$

Solutions

- Predictor Corrector Method(s) (= alternative integration rule)
- Proxy Simulation Scheme / Weak Scheme (discussed later)

Sensitivities in Monte Carlo

Partial Derivative with respect to Model Parameters

Monte Carlo Methods: Sensitivities

Let Z denote a random variable depending on realizations $Y := (X(T_1), \dots, X(T_m))$ of our simulated (Numéraire relative) state variables

$$Z = f(Y) = f(X(T_1), \dots, X(T_m))$$

e.g. the Numéraire relative path values of a financial product. Then the (Numéraire relative) price is given by

$$\mathbb{E}^{\mathbb{Q}}(Z \mid \mathcal{F}_{T_0}) = \mathbb{E}^{\mathbb{Q}}(f(X(T_1), \dots, X(T_m)) \mid \mathcal{F}_{T_0}).$$

Challenge: Let θ denote a parameter of the model SDE (e.g. its initial condition $X(0)$, volatility σ or any other complex function of those). We are interested in

$$\begin{aligned} \frac{\partial}{\partial \theta} \mathbb{E}^{\mathbb{Q}}(Z \mid \mathcal{F}_{T_0}) &= \frac{\partial}{\partial \theta} \int_{\Omega} f(X(T_1, \omega, \theta), \dots, X(T_m, \omega, \theta)) \, d\mathbb{Q}(\omega) \\ &= \frac{\partial}{\partial \theta} \int_{\mathbb{R}^m} \underbrace{f(x_1, \dots, x_m)}_{\substack{\text{payoff} \\ \text{may be} \\ \text{discontinuouse}}} \cdot \underbrace{\phi(X(T_1, \omega, \theta), \dots, X(T_m, \omega, \theta))(x_1, \dots, x_m)}_{\text{density - in general smooth in } \theta} \, d(x_1, \dots, x_m) \end{aligned}$$

Problem: Monte-Carlo approximation inherits regularity of f not of ϕ :

$$\mathbb{E}^{\mathbb{Q}}(Z \mid \mathcal{F}_{T_0}) \approx \hat{\mathbb{E}}^{\mathbb{Q}}(Z \mid \mathcal{F}_{T_0}) := \frac{1}{n} \sum_{i=1}^n \underbrace{f(X(T_1, \omega_i, \theta), \dots, X(T_m, \omega_i, \theta))}_{\substack{\text{payoff on path - may be} \\ \text{discontinuouse}}}$$

Monte Carlo Methods: Sensitivities

Finite Differences:

$$\begin{aligned}\frac{\partial}{\partial \theta} \mathbb{E}^{\mathbb{Q}}(f(Y(\theta)) \mid \mathcal{F}_{T_0}) &\approx \frac{\partial}{\partial \theta} \hat{\mathbb{E}}^{\mathbb{Q}}(f(Y(\theta)) \mid \mathcal{F}_{T_0}) \approx \frac{1}{2h} (\hat{\mathbb{E}}^{\mathbb{Q}}(f(Y(\theta + h)) \mid \mathcal{F}_{T_0}) - \hat{\mathbb{E}}^{\mathbb{Q}}(f(Y(\theta - h)) \mid \mathcal{F}_{T_0})) \\ &= \frac{1}{n} \sum_{i=1}^n \frac{1}{2h} (f(Y(\omega_i, \theta + h)) - f(Y(\omega_i, \theta - h)))\end{aligned}$$

Properties

- Requires no additional information from the model sde $dX = \dots$
- Requires no additional information from the simulation scheme $X(T_{i+1}) = \dots$
- Requires no additional information from the payout f
- Requires no additional information on the nature of θ ($\Rightarrow$ generic sensitivities)
- Biased derivative for *large* h due to finite difference of order h
- Extremely large variance for discontinuous payouts and *small* h (order h^{-1})

Monte Carlo Methods: Sensitivities: Pathwise Differentiation

Pathwise Differentiation:

$$\begin{aligned}\frac{\partial}{\partial \theta} \mathbb{E}^{\mathbb{Q}}(f(Y(\theta)) \mid \mathcal{F}_{T_0}) &\approx \frac{\partial}{\partial \theta} \hat{\mathbb{E}}^{\mathbb{Q}}(f(Y(\theta)) \mid \mathcal{F}_{T_0}) \\ &= \frac{1}{n} \sum_{i=1}^n \frac{\partial}{\partial \theta} (f(Y(\omega_i, \theta))) = \frac{1}{n} \sum_{i=1}^n f'(Y(\omega_i, \theta)) \cdot \frac{\partial Y(\omega_i, \theta)}{\partial \theta}\end{aligned}$$

Properties

- Requires additional information on the model sde $dX = \dots$
- Requires no additional information from the simulation scheme $X(T_{i+1}) = \dots$
- Requires additional information on the payout f (derivative of f must be known)
- Requires additional information on the nature of θ ($\Rightarrow$ restricted class of model parameters)
- Unbiased derivative
- Requires smoothness of payout? (in this formulation)

Monte Carlo Methods: Sensitivities: Pathwise Differentiation

Pathwise Differentiation (alternative interpretation):

$$\begin{aligned}\frac{\partial}{\partial \theta} \mathbb{E}^{\mathbb{Q}}(f(Y(\theta)) \mid \mathcal{F}_{T_0}) &= \frac{\partial}{\partial \theta} \int_{\Omega} f(Y(\omega, \theta)) \, d\mathbb{Q}(\omega) = \int_{\Omega} \frac{\partial}{\partial \theta} f(Y(\omega, \theta)) \, d\mathbb{Q}(\omega) \\ &= \int_{\Omega} f'(Y(\omega, \theta)) \cdot \frac{\partial Y(\omega, \theta)}{\partial \theta} \, d\mathbb{Q}(\omega) = \mathbb{E}^{\mathbb{Q}}(f'(Y(\theta)) \cdot \frac{\partial Y(\theta)}{\partial \theta} \mid \mathcal{F}_{T_0}) \\ &\approx \hat{\mathbb{E}}^{\mathbb{Q}}(f'(Y(\theta)) \cdot \frac{\partial Y(\theta)}{\partial \theta} \mid \mathcal{F}_{T_0}) = \frac{1}{n} \sum_{i=1}^n f'(Y(\omega_i, \theta)) \cdot \frac{\partial Y(\omega_i, \theta)}{\partial \theta}\end{aligned}$$

Note: See Joshi & Kainth [JK] or Rott & Fries [RF] for an example on how use pathwise differentiation with discontinuous payouts (there in the context of Default Swaps, CDOs).

Properties

- Requires additional information on the model sde $dX = \dots$
- Requires no additional information on the simulation scheme $X(T_{i+1}) = \dots$
- Requires additional information on the payout f (derivative of f must be known)
- Requires additional information on the nature of θ ($\Rightarrow$ restricted class of model parameters)
- Unbiased derivative
- Discontinuous payouts may be handled (interpret f' as distribution, for applications see e.g. [JK, RF])

Monte Carlo Methods: Sensitivities: Likelihood Ratio

Likelihood Ratio:

$$\begin{aligned}\frac{\partial}{\partial \theta} \mathbb{E}^{\mathbb{Q}}(f(Y(\theta)) \mid \mathcal{F}_{T_0}) &= \frac{\partial}{\partial \theta} \int_{\Omega} f(Y(\omega, \theta)) \, d\mathbb{Q}(\omega) = \frac{\partial}{\partial \theta} \int_{\mathbb{R}^m} f(y) \cdot \phi_{Y(\theta)}(y) \, dy \\ &= \int_{\mathbb{R}^m} f(y) \cdot \frac{\frac{\partial}{\partial \theta} \phi_{Y(\theta)}(y)}{\phi_{Y(\theta)}(y)} \cdot \phi_{Y(\theta)}(y) \, dy = \mathbb{E}^{\mathbb{Q}}(f(Y) \cdot w(\theta) \mid \mathcal{F}_{T_0}) \\ &\approx \hat{\mathbb{E}}^{\mathbb{Q}}(f(Y) \cdot w(\theta) \mid \mathcal{F}_{T_0}) = \frac{1}{n} \sum_{i=1}^n f(Y(\omega_i)) \cdot w(\theta, \omega_i)\end{aligned}$$

Properties

- Requires additional information on the model sde $dX = \dots (\rightarrow \phi_{Y(\theta)})$
- Requires no additional information on the simulation scheme $X(T_{i+1}) = \dots$
- Requires no additional information on the payout f
- Requires additional information on the nature of $\theta (\Rightarrow \text{restricted class of model parameters})$
- Unbiased derivative
- Discontinuous payouts may be handled.

Malliavin Calculus:

$$\begin{aligned}\frac{\partial}{\partial \theta} \mathbb{E}^{\mathbb{Q}}(f(Y(\theta)) \mid \mathcal{F}_{T_0}) &= \mathbb{E}^{\mathbb{Q}}(f(Y(\theta)) \cdot w(\theta) \mid \mathcal{F}_{T_0}) \\ &\approx \hat{\mathbb{E}}^{\mathbb{Q}}(f(Y(\theta)) \cdot w(\theta) \mid \mathcal{F}_{T_0}) = \frac{1}{n} \sum_{i=1}^n f(Y(\theta, \omega_i)) \cdot w(\theta, \omega_i)\end{aligned}$$

Note: Benhamou [B01] showed that the [Likelihood Ratio](#) corresponds to the Malliavin weights with minimal variance and may be expressed as a conditional expectation of all corresponding Malliavin weights (we thus view the Likelihood Ratio as an example for the Malliavin weighting method).

Properties

- Requires additional information on the model sde $dX = \dots (\rightarrow w)$
- Requires no additional information on the simulation scheme $X(T_{i+1}) = \dots$
- Requires no additional information on the payout f
- Requires additional information on the nature of $\theta (\Rightarrow \text{restricted class of model parameters})$
- Unbiased derivative
- Discontinuous payouts may be handled.

Proxy Simulation Scheme

Proxy Simulation Scheme

Pricing / Sensitivities

Proxy Scheme Simulation: Pricing

Proxy Scheme: Consider *three* stochastic processes

$X \quad t \mapsto X(t) \quad t \in \mathbb{R} \quad \text{model sde}$

$X^* \quad T_i \mapsto X^*(T_i) \quad i = 0, 1, 2, \dots \quad \text{time discretization scheme of } X \rightarrow \textit{target scheme}$

$X^\circ \quad T_i \mapsto X^\circ(T_i) \quad i = 0, 1, 2, \dots \quad \text{any other time discrete stochastic process}$
(assumed to be *close* to X^*) $\rightarrow$ *proxy scheme*

Pricing:

Let $Y = (X(T_1), \dots, X(T_m))$, $Y^* = (X^*(T_1), \dots, X^*(T_m))$, $Y^\circ = (X^\circ(T_1), \dots, X^\circ(T_m))$.

We have $E^{\mathbb{Q}}(f(Y(\theta)) \mid \mathcal{F}_{T_0}) \approx E^{\mathbb{Q}}(f(Y^*(\theta)) \mid \mathcal{F}_{T_0})$ and furthermore

$$\begin{aligned} E^{\mathbb{Q}}(f(Y^*(\theta)) \mid \mathcal{F}_{T_0}) &= \int_{\Omega} f(Y^*(\omega, \theta)) d\mathbb{Q}(\omega) = \int_{\mathbb{R}^m} f(y) \cdot \phi_{Y^*(\theta)}(y) dy \\ &= \int_{\mathbb{R}^m} f(y) \cdot \frac{\phi_{Y^*(\theta)}(y)}{\phi_{Y^\circ}(y)} \cdot \phi_{Y^\circ}(y) dy = E^{\mathbb{Q}}(f(Y^\circ) \cdot w(\theta) \mid \mathcal{F}_{T_0}) \end{aligned}$$

where $w(\theta) = \frac{\phi_{Y^*(\theta)}(y)}{\phi_{Y^\circ}(y)}$.

Note:

- For $X^\circ = X^*$ we have $w(\theta) = 1 \Rightarrow$ ordinary Monte Carlo.
- Y° is seen as being independent of θ . $\Rightarrow$ implications on sensitivities.
- Requirement: $\forall y : \phi^{Y^\circ}(y) = 0 \Rightarrow \phi^{Y^*}(y) = 0$

Proxy Scheme Simulation: Sensitivities

Proxy Scheme Sensitivities:

$$\begin{aligned}\frac{\partial}{\partial \theta} \mathbb{E}^{\mathbb{Q}}(f(Y^*(\theta)) \mid \mathcal{F}_{T_0}) &= \frac{\partial}{\partial \theta} \int_{\Omega} f(Y^*(\omega, \theta)) \, d\mathbb{Q}(\omega) = \frac{\partial}{\partial \theta} \int_{\mathbb{R}^m} f(y) \cdot \phi_{Y^*(\theta)}(y) \, dy \\ &= \int_{\mathbb{R}^m} f(y) \cdot \frac{\frac{\partial}{\partial \theta} \phi_{Y^*(\theta)}(y)}{\phi_{Y^\circ}(y)} \cdot \phi_{Y^\circ}(y) \, dy = \mathbb{E}^{\mathbb{Q}}(f(Y^\circ) \cdot w'(\theta) \mid \mathcal{F}_{T_0}) \\ &\approx \hat{\mathbb{E}}^{\mathbb{Q}}(f(Y^\circ) \cdot w'(\theta) \mid \mathcal{F}_{T_0}) = \frac{1}{n} \sum_{i=1}^n f(Y^\circ(\omega_i)) \cdot w'(\theta, \omega_i)\end{aligned}$$

Properties

- Requires no additional information on the model sde $dX = \dots$
- Requires additional information on the simulation scheme $X^*(T_{i+1}), X^\circ(T_{i+1})$
- Requires no additional information on the payout f
- Requires additional information on the nature of θ ($\Rightarrow$ restricted class of model parameters)
- Unbiased derivative (biased if finite differences are used for w)
- Discontinuous payouts may be handled.

Proxy Scheme Simulation: Sensitivities

Proxy Scheme Sensitivities:

$$\begin{aligned}\frac{\partial}{\partial \theta} \mathbb{E}^{\mathbb{Q}}(f(Y^*(\theta)) \mid \mathcal{F}_{T_0}) &\approx \frac{1}{2h} (\mathbb{E}^{\mathbb{Q}}(f(Y^*(\theta + h)) \mid \mathcal{F}_{T_0}) - \mathbb{E}^{\mathbb{Q}}(f(Y^*(\theta - h)) \mid \mathcal{F}_{T_0})) \\ &= \frac{\partial}{\partial \theta} \int_{\mathbb{R}^m} f(y) \cdot \frac{1}{2h} (\phi_{Y^*(\theta+h)}(y) - \phi_{Y^*(\theta-h)}(y)) \, dy \\ &= \int_{\mathbb{R}^m} f(y) \cdot \frac{\frac{1}{2h} (\phi_{Y^*(\theta+h)}(y) - \phi_{Y^*(\theta-h)}(y))}{\phi_{Y^\circ}(y)} \cdot \phi_{Y^\circ}(y) \, dy \\ &\approx \frac{1}{n} \sum_{i=1}^n f(Y^\circ(\omega_i)) \cdot \frac{1}{2h} (w(\theta + h, \omega_i) - w(\theta - h, \omega_i))\end{aligned}$$

Properties

- Requires no additional information on the model sde $dX = \dots$
- Requires additional information on the simulation scheme $X^*(T_{i+1}), X^\circ(T_{i+1})$
- Requires no additional information on the payout f
- Requires no additional information on the nature of θ ($\Rightarrow$ generic sensitivities)
- Biased derivative (but small shift h possible!)
- Discontinuous payouts may be handled.

Proxy Scheme Simulation: Sensitivities

Proxy Scheme Sensitivities:

$$\begin{aligned}\frac{\partial}{\partial \theta} \mathbb{E}^{\mathbb{Q}}(f(Y^*(\theta)) \mid \mathcal{F}_{T_0}) &\approx \frac{1}{2h} (\mathbb{E}^{\mathbb{Q}}(f(Y^*(\theta + h)) \mid \mathcal{F}_{T_0}) - \mathbb{E}^{\mathbb{Q}}(f(Y^*(\theta - h)) \mid \mathcal{F}_{T_0})) \\ &= \frac{\partial}{\partial \theta} \int_{\mathbb{R}^m} f(y) \cdot \frac{1}{2h} (\phi_{Y^*(\theta+h)}(y) - \phi_{Y^*(\theta-h)}(y)) \, dy \\ &= \int_{\mathbb{R}^m} f(y) \cdot \frac{\frac{1}{2h} (\phi_{Y^*(\theta+h)}(y) - \phi_{Y^*(\theta-h)}(y))}{\phi_{Y^\circ}(y)} \cdot \phi_{Y^\circ}(y) \, dy \\ &\approx \frac{1}{n} \sum_{i=1}^n f(Y^\circ(\omega_i)) \cdot \frac{1}{2h} (w(\theta + h, \omega_i) - w(\theta - h, \omega_i))\end{aligned}$$

*Finite difference applied to the pricing
results in a finite difference approximation of the Likelihood Ratio*

thus

We have all the nice properties of the *Likelihood Ratio*
combined with the genericity of *Finite Differences*

Proxy Simulation Scheme

Densities / Weak Schemes

Proxy Scheme Simulation: Densities / Weak Schemes

Proxy Scheme: Consider *three* stochastic processes

$X \quad t \mapsto X(t) \quad t \in \mathbb{R} \quad \text{model sde}$

$X^* \quad T_i \mapsto X^*(T_i) \quad i = 0, 1, 2, \dots \quad \text{time discretization scheme of } X \rightarrow \text{target scheme}$

$X^\circ \quad T_i \mapsto X^\circ(T_i) \quad i = 0, 1, 2, \dots \quad \text{any other time discrete stochastic process}$
(assumed to be *close* to X^*) $\rightarrow$ *proxy scheme*

Pricing: Let $Y = (X(T_1), \dots, X(T_m))$, $Y^* = (X^*(T_1), \dots, X^*(T_m))$, $Y^\circ = (X^\circ(T_1), \dots, X^\circ(T_m))$.

$$\mathbb{E}^{\mathbb{Q}}(f(Y(\theta)) \mid \mathcal{F}_{T_0}) \approx \mathbb{E}^{\mathbb{Q}}(f(Y^*(\theta)) \mid \mathcal{F}_{T_0}) = \mathbb{E}^{\mathbb{Q}}(f(Y^\circ) \cdot w(\theta) \mid \mathcal{F}_{T_0})$$

where


$$w(\theta) = \frac{\phi_{Y^*(\theta)}(y)}{\phi_{Y^\circ}(y)} \quad (\text{calculated numerically}).$$

Note:


- From the scheme X° we need the realizations (to generate the path)
 $\rightarrow$ Need something explicit (Euler-Scheme, Predictor Corrector, etc.)
- From the scheme X^* we need the transition probability only (weaker requirement)
 $\rightarrow$ May use complex implicit schemes or expansions of the the transition probability of the (true) model sde.
Kampen derived a quadratic WKB expansion for the LIBOR Market Model (see appendix)

Proxy Simulation: Implementation


Standard Monte Carlo Simulation: Pricing




Standard Monte Carlo Simulation: Sensitivities



Proxy Simulation Method: Pricing



Proxy Simulation Method: Sensitivities



LR like Sensitivity as
Finite Difference

Example: LIBOR Market Model

Example: Proxy Scheme Simulation for a LIBOR Market Model

LIBOR Market Model:

$$dL_i = L_i \mu_i^L dt + L_i \sigma_i dW_i, \quad i = 1, \dots, n, \quad \text{with} \quad \mu_i^L = \sum_{i < j \leq n} \frac{L_j \delta_j}{1 + L_j \delta_j} \sigma_i \sigma_j \rho_{i,j}, \quad dW = \Sigma \cdot \Gamma \cdot dU,$$

where $dW = (dW_1, \dots, dW_n)$, $dW_i dW_j = \rho_{i,j} dt$, $\Sigma = \text{diag}(\sigma_1, \dots, \sigma_n)$, $\Gamma \Gamma^T = (\rho_{i,j})$.

- Log-normal model (common extensions: local vol., stoch. vol., jump)
- Non-linear drift
- High dimensional (no low dimensional Markovian state variable)
- Driving factors may be low dimensional (parsimonious model) $\rightarrow \Gamma$ is an $n \times m$ matrix.

LIBOR Market Model & Numerical Schemes in Log-Coordinates:

$$\text{model sde: } dK = \mu^K dt + \Sigma \cdot \Gamma \cdot dU \quad K := \log(L), \quad \mu^K := \mu^L - \frac{1}{2} \Sigma^2$$

$$\text{proxy scheme: } K^\circ(T_{i+1}) = K^\circ(T_i) + \mu^{K^\circ}(T_i) \Delta T_i + \Sigma^\circ(T_i) \cdot \Gamma^\circ(T_i) \cdot \Delta U(T_i)$$

$$\text{target scheme: } K^*(T_{i+1}) = K^*(T_i) + \mu^{K^*}(T_i) \Delta T_i + \Sigma(T_i) \cdot \Gamma(T_i) \cdot \Delta U(T_i)$$

Example: Proxy Scheme Simulation for a LIBOR Market Model

LIBOR Market Model & Numerical Schemes in Log-Coordinates:

$$\text{model sde: } dK = \mu^K dt + \Sigma \cdot \Gamma \cdot dU \quad K := \log(L), \quad \mu^K := \mu^L - \frac{1}{2}\Sigma^2$$

$$\text{proxy scheme: } K^\circ(T_{i+1}) = K^\circ(T_i) + \mu^{K^\circ}(T_i)\Delta T_i + \Sigma^\circ(T_i) \cdot \Gamma^\circ(T_i) \cdot \Delta U(T_i) \quad \leftarrow \text{sample path}$$

$$\text{target scheme: } K^*(T_{i+1}) = K^*(T_i) + \mu^{K^*}(T_i)\Delta T_i + \Sigma(T_i) \cdot \Gamma(T_i) \cdot \Delta U(T_i)$$

Transition Probabilities $T_i \rightarrow T_{i+1}$:

Assume for simplicity that $\mu^{K^*}(T_i)$ depends on $K^*(T_i)$, $K^*(T_{i+1})$ only (and same for $^\circ$)
($\rightarrow$ true for, e.g. Euler Scheme, Predictor Corrector), then

$$\phi^{K^\circ}(T_i, K_i^\circ; T_{i+1}, K_{i+1}^\circ) = \frac{1}{(2\pi\Delta T_i)^{n/2}} \exp\left(-\frac{1}{2\Delta T_i} (\Lambda^{\circ-1/2} F^{\circ T} \Sigma^{\circ-1} (K_{i+1}^\circ - K_i^\circ - \mu^{K^\circ}(T_i)\Delta T_i))^2\right)$$

$$\phi^{K^*}(T_i, K_i^*; T_{i+1}, K_{i+1}^*) = \frac{1}{(2\pi\Delta T_i)^{n/2}} \exp\left(-\frac{1}{2\Delta T_i} (\Lambda^{-1/2} F^T \Sigma^{-1} (K_{i+1}^* - K_i^* - \mu^{K^*}(T_i)\Delta T_i))^2\right)$$

Proxy Scheme Weights:

$$w(T_{i+1}) | \mathcal{F}_{T_k} = \prod_{j=k}^i \frac{\phi^{K^*}(T_j, K_j^\circ; T_{j+1}, K_{j+1}^\circ)}{\phi^{K^\circ}(T_j, K_j^\circ; T_{j+1}, K_{j+1}^\circ)} \quad \leftarrow \text{monte carlo weights}$$

Note: We used the factor decomposition (PCA) $\Gamma = F \cdot \sqrt{\Lambda}$ where $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_m)$ are the non-zero Eigenvalues of $\Gamma \cdot \Gamma^T$.

A **change of market data / calibration** enters into transition probabilities only.

Summary: Requirements / Implementation

Proxy Scheme Weights:

$$w(T_{i+1}) |_{\mathcal{F}_{T_k}} = \prod_{j=k}^i \frac{\phi^{K^*}(T_j, K_j^\circ; T_{j+1}, K_{j+1}^\circ)}{\phi^{K^\circ}(T_j, K_j^\circ; T_{j+1}, K_{j+1}^\circ)}$$

Implementation:

The transition densities ϕ^{K° and ϕ^{K^} are densities from the numerical schemes K° and K^* . They may be calculated numerically (on the fly together with the (proxy) schemes paths)!*

Requirement:

$$\phi^{K^\circ}(T_i, K_i^\circ; T_{i+1}, K_{i+1}^\circ) = 0 \implies \phi^{K^*}(T_i, K_i^\circ; T_{i+1}, K_{i+1}^\circ) = 0$$

This requirement corresponds to the non-degeneracy condition imposed on the diffusion matrix in the continuous case (e.g. Malliavin Calculus).

However: Here, this requirement may be achieved even for a degenerate diffusion matrix, e.g. by a non-linear drift.

Moreover:

Since we are free to choose the proxy scheme, it may be chosen such that the condition holds.

Summary: Properties / Achievements

Properties:

- Requires no additional information on the model sde $dX = \dots$
- Requires additional information on the simulation scheme $X^*(T_{i+1}), X^\circ(T_{i+1})$
- Requires no additional information on the payout f
- Requires no additional information on the nature of θ ($\Rightarrow$ generic sensitivities)
- Stable for small shifts h
- Discontinuous payouts may be handled.

Achievements:

- **Stable Generic Sensitivites:** Finite Differences result in numerical Likelihood Ratios
- **Weak Schemes:** Allows to correct for an improper transition density.

Summary: Note on the non-degeneracy condition


A note on the requirement

$$\forall y : \phi^{Y^\circ}(y) = 0 \Rightarrow \phi^{Y^*}(y) = 0$$

While for Malliavin Calculus one would expect some non-degeneracy condition imposed on the diffusion matrix, the above may hold for (time-discrete) simulation schemes with degenerate diffusion matrices, e.g. if a stochastic drift generates additional diffusion.

Example:

Consider a model on two state variable (here an LMM) with a degenerate (rank 1) diffusion matrix (**red**) and a stochastic drift term (like in LMM). Then a single Euler step will span a line (**blue**). Using this as a proxy scheme will not allow drift corrections outside that 1-dim hypersurface. However, two subsequent Euler steps of half the size, generate diffusion perpendicular to the 1-dim hypersurface (**green**).



Examples and Numerical Results

Numerical Results

Proxy Scheme: Consider *three* stochastic processes

X $t \mapsto X(t)$ $t \in \mathbb{R}$ model sde

X^* $T_i \mapsto X^*(T_i)$ $i = 0, 1, 2, \dots$ time discretization scheme of $X \rightarrow$ *target scheme*

X° $T_i \mapsto X^\circ(T_i)$ $i = 0, 1, 2, \dots$ any other time discrete stochastic process
(assumed to be *close* to X^*) $\rightarrow$ *proxy scheme*

Test Case:

X LIBOR Market Model

X^* Target Scheme: Some standard discretization of LMM.


X° Proxy Scheme: Log-normal scheme without drift (LMM drift zero) ([extrem test case](#)).

Check for:

- Bond prices ($\Leftrightarrow$ can we correct for the drift)
- Sensitivities of Trigger Products (Digitals, Auto Caps)


Example I: Correcting the Drift

Numerical Results: Monte Carlo Bond Price Distributions




- Shown: Absolute Bond price Monte Carlo error distribution for Euler Scheme with drift zero (red) and Euler Scheme with Euler drift (yellow): Neglecting drift results in large Bond price errors and even higher Monte Carlo variance (since here drift would generate mean reversion).
- Next: Use zero-drift Euler Scheme as proxy scheme and correct drift towards Euler Scheme with drift (target scheme).

Numerical Results: Monte Carlo Bond Price Distributions




- Shown: Use zero-drift Euler Scheme as proxy scheme (red) and correct drift towards Euler Scheme with drift (target scheme, blue)
- Next: Take a closer look. Compare proxy scheme simulation with direct simulation

Numerical Results: Monte Carlo Bond Price Distributions



- Shown: Monte Carlo Error of Bond Prices for Proxy-Scheme Method (using zero-drift Euler Scheme as proxy scheme) (blue) and direct simulation of target scheme (yellow)
- Next: Refine target scheme by more accurate transition probabilities

Numerical Results: Monte Carlo Bond Price Distributions




- Shown: Direct Euler Scheme simulation (yellow), Proxy Scheme simulation with Euler Scheme as target scheme (blue), Proxy Scheme simulation with transition probabilities derived from trapezoidal integration rule for the drift (green).

Example 2:


Robust Generic Sensitivities

Numerical Results: Monte Carlo Sensitivities




- Proxy Scheme Sensitivity shows an increase of variance for large shift (well known effect for Likelihood Ratio / Malliavin Calculus)
- Proxy Scheme Sensitivity remains stable for small shifts

Numerical Results: Monte Carlo Sensitivities




- Proxy Scheme Sensitivity shows an increase of variance for large shift (well known effect for Likelihood Ratio / Malliavin Calculus)
- Proxy Scheme Sensitivity remains stable for small shifts

Numerical Results: Monte Carlo Sensitivities



- Proxy Scheme Sensitivity shows an increase of variance for large shift (well known effect for Likelihood Ratio / Malliavin Calculus)
- Proxy Scheme Sensitivity remains stable for small shifts

Numerical Results: Monte Carlo Sensitivities



- Proxy Scheme Sensitivity shows an increase of variance for large shift (well known effect for Likelihood Ratio / Malliavin Calculus)
- Proxy Scheme Sensitivity remains stable for small shifts

Appendix

Appendix: Quadratic WKB Expansion for the LMM Transition Probability Density

Three assumptions. First

- (A) The operator L is uniformly parabolic in $\mathbb{R}^n$, i.e. there exists $0 < \lambda < \Lambda < \infty$ such that for all $\xi \in \mathbb{R}^n \setminus \{0\}$

$$0 < \lambda \leq \sum_{i,j=1}^n a_{ij}(x) \xi_i \xi_j \leq \Lambda. \quad (1)$$

- (B) The coefficients of L are bounded functions in $\mathbb{R}^n$ which are uniformly Hölder continuous of exponent α ($\alpha \in (0, 1)$).

guarantee that fundamental solution exists and is strictly positive. The third assumption

- (C) the growth of all derivatives of the smooth coefficients functions $x \rightarrow a_{ij}(x)$ and $x \rightarrow b_i(x)$ is at most of exponential order, i.e. there exists for each multiindex α a constant $\lambda_\alpha > 0$ such that for all $1 \leq i, j, k \leq n$

$$\left| \frac{\partial^\alpha a_{jk}}{\partial x^\alpha} \right|, \left| \frac{\partial^\alpha b_i}{\partial x^\alpha} \right| \leq \exp(\lambda_\alpha |x|^2), \quad (2)$$

guarantees (pointwise) convergence of coefficient functions $x \rightarrow c_k^y(x) := c_k(x, y)$ and $x \rightarrow d_k^y(x) := d_k(x, y)$ in the standard WKB-expansion

$$p(\delta t, x, y) = \frac{1}{\sqrt{2\pi\delta t}^n} \exp \left(-\frac{d^2(x, y)}{2\delta t} + \sum_{i \geq 0} c_i(x, y) \delta t^i \right). \quad (3)$$

and in the new WKB expansion (we call it the quadratic WKB expansion), which is

$$p(\delta t, x, y) = \frac{1}{\sqrt{2\pi\delta t}^n} \times \exp \left(-\frac{(\sum_{i \geq 0} d_i \delta t^i)^2}{2\delta t} + \sum_{i \geq 0} (c_i^y(y) + \nabla \left(c_i^y - \sum_{l=1}^{i-1} d_l^y d_{i-l}^y \right) (y) \cdot (x - y)) \delta t^i \right). \quad (4)$$

(This is from the ansatz

$$p(\delta t, x, y) = \frac{1}{\sqrt{2\pi\delta t}^n} \exp \left(-\frac{(\sum_{i \geq 0} d_i \delta t^i)^2}{2\delta t} + \sum_{i \geq 0} (\alpha_i^y + \beta_i^y \cdot (x - y)) \delta t^i \right). \quad (5)$$

where $\cdot$ denotes the scalar product. Here α_i^y and β_i^y are affine terms depending on y (compensation terms).

From the target scheme only the transition probability is needed.

Kampen [KF] derived a quadratic WKB expansion for the LIBOR Market Model (see left).

This enables us to construct a proxy scheme simulation with almost arbitrary small time discretization error - even for a large time steps ΔT .

References

References

- [B01] BENHAMOU, ERIC: Optimal Malliavin Weighting Function for the Computation of the Greeks. 2001.
- [BG96] BROADIE, MARK; GLASSERMAN, PAUL: Estimating Security Price Derivatives using Simulation. *Management Science*, 1996, Vol. 42, No. 2, 269-285.
- [F] FRIES, CHRISTIAN P.: Bumping the Model. 2005.
<http://www.christian-fries.de/finmath/bumpingthemodel>
- [FK] FRIES, CHRISTIAN P.; KAMPEN, JÖRG: Proxy Simulation Schemes for generic robust Monte-Carlo sensitivities and high accuracy drift approximation (with applications to the LIBOR Market Model). 2005.
<http://www.christian-fries.de/finmath/proxyscheme>
- [FLLLT] FOURNIÉ, ERIC; LASRY JEAN-MICHEL; LEBUCHOUX, JÉRÔME; LIONS, PIERRE-LOUIS; TOUZI, NIZAR: Applications of Malliavin calculus to Monte Carlo methods in finance. *Finance Stochastics*. 3, 391-412 (1999). Springer- Verlag 1999.
- [G03] GLASSERMAN, PAUL: Monte Carlo Methods in Financial Engineering. (Stochastic Modelling and Applied Probability). Springer, 2003. ISBN 0-387-00451-3.
- [JK] JOSHI, MARK S.; KAINTH, DHERMINDER: Rapid computation of prices and deltas of n^{th} to default swaps in the Li Model. *Quantitative Finance*, volume 4, issue 3, (June 04), p. 266- 275.
<http://www.quarchome.org/>.
- [KF] KAMPEN, JÖRG; FRIES, CHRISTIAN: A Quadratic WKB Expansion for the Transition Probability of the LIBOR Market Model. *in preparation*
- [M97] MALLIAVIN, PAUL: Stochastic Analysis (Grundlehren Der Mathematischen Wissenschaften). Springer Verlag, 1997. ISBN 3-540-57024-1 .
- [RF] ROTT, MARIUS G.; FRIES, CHRISTIAN P.: Fast and Robust Monte Carlo CDO Sensitivities and their Efficient Object Oriented Implementation. 2005.
<http://www.christian-fries.de/finmath/cdogreeks>